

REPORT DOCUMENTATION PAGE				<i>Form Approved OMB No. 0704-0188</i>	
<small>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</small>					
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)

PUBLICATION OR PRESENTATION RELEASE REQUEST

Pubkey: 9229

NRLINST 5600.2

Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D	() Abstract only, published () Book () Conference Proceedings (refereed) () Invited speaker () Journal article (refereed) () Oral Presentation, published () Other, explain	() Abstract only, not published () Book chapter () Conference Proceedings (not refereed) () Multimedia report (X) Journal article (not refereed) () Oral Presentation, not published	STRN <u>NRLJA/7320-14-2170</u> Route Sheet No. <u>7320/</u> Job Order No. <u>73-8677-03-5</u> Classification <u>X</u> U C Sponsor <u>ONR</u> approval obtained <u>yes</u> <u>X</u> no
Encl: (1) Two copies of subject paper (or abstract)			

Title of Paper or Presentation

Inserting Tides and Topographic Wave Drag into High-resolution Eddyding Simulations

Author(s) Name(s) (First, MI, Last), Code, Affiliation if not NRL

Brian Arbic Univ. of Michigan Maarten Buijsman University of New Orleans Eric Chassignet Florida State University S.T. Garner NOAA - New Jersey
John A. Goff Univ. of Texas at Austin S.R. Jayne WHOI E. Joseph Metzger 7323 James G Richman 7323 Jay Shriver 7323
Patrick G. Timko Bangor Univ. - UK D.S. Trossman Univ. of Michigan Alan J. Wallcraft 7323

It is intended to offer this paper to the

(Name of Conference)

and/or for publication in CLIVAR Exchanges Newsletter (Date, Place and Classification of Conference)
Exchanges - CLIVAR Newsletter, Unclassified (Name and Classification of Publication)

(Name of Publisher)

After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a).

It is the opinion of the author that the subject paper (is) (is not X) classified, in accordance with reference (b).

This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does) (does not X) contain any militarily critical technology.
This subject paper (has) (has never X) been incorporated in an official NRL Report.

E. Joseph Metzger, 7323

Name and Code (Principal Author)

E. Joseph Metzger
(Signature)

CODE	SIGNATURE	DATE	COMMENTS
Author(s) <u>Metzger</u>	<i>E. Joseph Metzger</i>	<u>5/12/2014</u>	Need by <u>16 Jun 14</u> Publicly accessible sources used for this publication
Section Head <u>Hogan</u>	<i>Patricia Hogan</i>	<u>5/14/14</u>	
Branch Head Gregg A. Jacobs, 7320	<i>Gregg A. Jacobs</i>	<u>5/13/14</u>	
Division Head Ruth H. Preller, 7300	<i>Ruth H. Preller</i>	<u>5/23/14</u>	1. Release of this paper is approved. 2. To the best knowledge of this Division, the subject matter of this paper (has <u> </u>) (has never <u>X</u>) been classified.
Security, Code 1231			1. Paper or abstract was released. 2. A copy is filed in this office.
Office of Counsel, Code 1008.3	<i>Kathy Chapman</i>	<u>6/3/14</u>	
ADOR/Director NCST E. R. Franchi, 7000			
Public Affairs (Unclassified/ Unlimited Only), Code 7030.4	<i>Shannon Menneri</i>	<u>6-2-14</u>	
Division, Code			
Author, Code			

PUBLICATION OR PRESENTATION RELEASE REQUEST

14-1231-1735

Publ 9229

NRLINST 5600.2

Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D	() Abstract only, published () Book () Conference Proceedings (refereed) () Invited speaker () Journal article (refereed) () Oral Presentation, published () Other, explain	() Abstract only, not published () Book chapter () Conference Proceedings (not refereed) () Multimedia report (X) Journal article (not refereed) () Oral Presentation, not published	STRN <u>NRL/IA/7320-14-2170</u> Route Sheet No. <u>7320</u> Job Order No. <u>73-8677-03-5</u> Classification <u>SECRET</u> Sponsor <u>ONR</u> approval obtained <u>yes</u> <u>X</u> no
---	--	--	---

Title of Paper or Presentation

Inserting Tides and Topographic Wave Drag into High-resolution Eddyling Simulations

Author(s) Name(s) (First, M.I., Last), Code, Affiliation if not NRL

Brian Arlic Univ. of Michigan Maarten Bulsman University of New Orleans Eric Chassignet Florida State University S.T. Garner NOAA - New Jersey
John A. Goff Univ. of Texas at Austin S.R. Jayne WHOI E. Joseph Metzger 7323 James G Richman 7323 Jay Shriver 7323
Patrick G. Timko Bangor Univ. - UK D.S. Tressman Univ. of Michigan Alan J. Wallcraft 7323

It is intended to offer this paper to the

(Name of Conference)

(Date, Place and Classification of Conference)

and/or for publication in

CLIVAR Exchanges Newsletter
Exchanges - CLIVAR Newsletter, Unclassified
(Name and Classification of Publication)

(Name of Publisher)

After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a).

It is the opinion of the author that the subject paper (is) (is not X) classified, in accordance with reference (b).This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does) (does not X) contain any militarily critical technology.This subject paper (has) (has never X) been incorporated in an official NRL Report.

E. Joseph Metzger, 7323

Name and Code (Principal Author)

E. Joseph Metzger
(Signature)



CODE	SIGNATURE	DATE	COMMENTS
Author(s) <u>Metzger</u>	<i>E. Joseph Metzger</i>	5/12/2014	Need by <u>16 Jun 14</u> Publicly accessible sources used for this publication This is a Final Security Review. Any changes made in the document, after approved by Code 1231, nullify the Security Review.
Section Head <u>Hogan</u>	<i>Patrick Hogan</i>	5/14/14	
Branch Head <u>Gregg A. Jacobs, 7320</u>	<i>Gregg A. Jacobs</i>	5/13/14	
Division Head <u>Ruth H. Preller, 7300</u>	<i>Ruth H. Preller</i>	5/23/14	1. Release of this paper is approved. 2. To the best knowledge of this Division, the subject matter of this paper (has <u> </u>) (has never <u>X</u>) been classified.
Security, Code <u>1231</u>	<i>Shannon Menni</i>	5/27/14	1. Paper or abstract was released. 2. A copy is filed in this office.
Office of Counsel, Code <u>1008.3</u>	<i>Kathy Chapman</i>	6/3/14	
ADOR/Director NCST <u>E. R. Franchi, 7000</u>			
Public Affairs (Unclassified/Unlimited Only), Code <u>7030.4</u>	<i>Shannon Menni</i>	6-2-14	
Division, Code			
Author, Code			



Exchanges

No.65 (Vol 19 No.2) July 2014

Special Issue: High Resolution Ocean Climate Modelling



CLIVAR Ocean & Climate: Variability, Predictability and Change is the World Climate Research Programme's (WCRP) project on ocean-atmosphere interactions.

WCRP is sponsored by the World Meteorological Organization, the International Council for Science and the Intergovernmental Oceanographic Commission of UNESCO.

WCRP
World Climate Research Programme



ICSU
International Council for Science

Inserting tides and topographic wave drag into high-resolution eddy simulations

Brian K. Arbic^{1*}, Maarten C. Buijsman², Eric P. Chassignet³, Stephen T. Garner⁴, Steven R. Jayne⁵, E. Joseph Metzger⁶, James G. Richman⁶, Jay F. Shriver⁶, Patrick G. Timko^{1,7}, David S. Trossman^{1,8}, and Alan J. Wallcraft⁶

1 Department of Earth and Environmental Sciences, University of Michigan, Ann Arbor, Michigan, USA

2 Department of Physics, University of New Orleans, New Orleans, Louisiana, USA

3 Department of Earth, Ocean, and Atmospheric Science and Center for Ocean-Atmospheric Prediction Studies, Florida State University, Tallahassee, Florida, USA

4 National Oceanic and Atmospheric Administration (NOAA)/Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey, USA

5 Physical Oceanography Department, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, USA

6 Oceanography Division, Naval Research Laboratory (NRL-SSC), Stennis Space Center, Mississippi, USA

7 Center for Applied Marine Sciences, Marine Science Laboratories, Bangor University, Menai Bridge, Anglesey, UK

8 Department of Atmospheric and Oceanic Sciences, McGill University, Montreal, Canada

Corresponding author: arbic@umich.edu

1. Introduction

We report here on two of our recent development efforts for global high-resolution ocean models, undertaken in the HYbrid Coordinate Ocean Model (HYCOM; Chassignet et al., 2009). The simulations discussed here are “eddy” (Hecht and Hasumi, 2008), meaning that they include an energetic mesoscale eddy field. In one of our efforts (Arbic

et al., 2010, 2012; Timko et al. 2012, 2013; Shriver et al. 2012, 2014; Richman et al., 2012), we have inserted tides into global eddy simulations. Tidal flows in a high-resolution global model with realistic rough topography generate internal tides-internal waves of tidal frequency. Similarly, geostrophic flows over rough topography generate internal lee waves (Bell, 1975). It is not yet possible to resolve internal lee waves in global models, but the impact of internal lee waves can be parameterised. In the other effort reported here, we have inserted parameterised topographic lee wave drag into global eddy simulations that do not include tides, with a preliminary focus on the resulting model energy budget (Trossman et al., 2013).

Global simulations of the tides and the atmospherically forced eddy general circulation have long been done separately. Global general circulation models began to resolve mesoscale eddies in the 1990’s (e.g., McClean et al., 1997). Since then eddy models have become increasingly realistic (e.g., Hecht and Hasumi, 2008). Global modelling of internal waves is a newer endeavour. In the first global simulations of internal tides (Arbic et al., 2004; Simmons et al., 2004), only tidal forcing was employed, and the stratification was taken to be horizontally uniform. The insertion of tides into atmospherically forced general circulation models allows for internal tide propagation in a realistic horizontally varying stratification, for interactions between internal tides and eddies, and for the co-existence of tides and near-inertial waves, another important class of internal waves (e.g., Simmons and Alford, 2012). In global high-resolution models with simultaneous atmospheric and tidal forcing, the internal wave spectrum is beginning to be resolved, just as mesoscale eddies began to be resolved in global models two decades ago. Our simulations with simultaneous tidal and atmospheric forcing are being used to address a variety of scientific and operational questions.

The global energy budget has been a topic of great recent interest, largely because the mixing occurring when energy is dissipated may exert a strong control on the oceanic meridional overturning circulation (Munk and Wunsch, 1998). The mechanisms underlying the dissipation of the eddy oceanic general circulation are still under investigation. Candidate dissipation mechanisms include the transfer of energy from geostrophic flows to submesoscale motions in the upper ocean (e.g., Capet et al., 2008), bottom boundary layer drag (Sen et al., 2008; Arbic et al., 2009), and breaking of internal lee waves generated by geostrophic flows over rough topography (e.g., Naveira-Garabato et al., 2004). Nikurashin and Ferrari (2011) and Scott et al. (2011) estimated the globally integrated energy flux of geostrophic flows into internal lee waves over rough topography, and found it to be a substantial fraction of the ~1 TW of energy put by the wind into geostrophic flows (e.g., Wunsch, 1998, and others). Both the Nikurashin and Ferrari (2011) and Scott et al. (2011) estimates were “offline”; they utilised bottom flows from eddy simulations that did not employ topographic wave drag as they ran. In Trossman et al. (2013) we inserted inline topographic lee wave drag into eddy HYCOM simulations, thus ensuring feedbacks between the model bottom stratification and flow fields and the topographic internal lee wave drag.

The HYCOM simulations shown here are forced by output from the Navy Operational Global Atmospheric Prediction System (NOGAPS); going forward the simulations will be forced by the Navy Global Environmental Model (NAVGEN). The simulations shown here are run on a combination Mercator and tripolar grid with an equatorial grid resolution of 0.08° (8.9 km) and are coupled to a sea ice model at high

latitudes. Global HYCOM simulations at 0.04° (4.5 km) have also been performed, but are not discussed here (see article by Chassignet et al. 2014, this issue). The model reproduces the general circulation of the global ocean, the strength and variability of the western boundary currents, such as the Gulf Stream and Kuroshio, and the mesoscale eddies generated by instabilities of the major currents (Thoppil et al., 2011).

2. Insertion of tides into eddying simulations

In our HYCOM simulations forced by both atmospheric fields and tides (Arbic et al., 2010, 2012), we include the four largest semidiurnal tidal constituents (M2, S2, N2, and K2) and the four largest diurnal tidal constituents (K1, O1, P1, and Q1). We use the parameterised topographic internal wave drag scheme of Garner (2005), modified to limit the maximum decay rates to (9 hours)⁻¹. The wave drag parameterisation represents the drag resulting from the generation and subsequent breaking of unresolved small-vertical scale internal waves by tidal flow over rough topography. Because the wave drags acting on tidal versus non-tidal motions have different strengths (Bell, 1975), tidal and non-tidal bottom flows are separated using a 25-hour running boxcar filter. Presently, the scalar approximation (Ray, 1998) is used for the self-attraction and loading term (Hendershott, 1972).

Barotropic tidal sea surface elevations in HYCOM have been compared to measurements made from tide gauges (Shum et al., 1997) and state-of-the-art data-assimilative barotropic tide models (Egbert et al., 1994). In Shriver et al. (2012) we also compared the modelled internal tide perturbation to sea surface height (SSH) - computed from a spatial high-pass of the total amplitude of tidal SSH - to internal tide perturbations computed from along-track satellite altimeter

data (Ray and Byrne, 2010). Figure 1 shows global maps of the M2 internal tide amplitude in HYCOM versus along-track altimeter data. The hotspots around locations such as Hawai'i, the Aleutian Islands, the Tuamotu Archipelago in the tropical central South Pacific, and Madagascar match up reasonably well in the two maps. Furthermore, the root-mean-square (rms) perturbation magnitudes over the five hotspot regions delineated by boxes in the bottom panel of Figure 1 agree with the altimeter rms values to within about 20%. Comparisons of tidal currents, and their vertical structure, in HYCOM versus historical moored observations were made in Timko et al. (2012; 2013).

We have used the HYCOM tidal simulations to distinguish between tidal and non-tidal contributions to the wavenumber spectrum of SSH (Richman et al., 2012). The slope of the wavenumber spectrum is of great theoretical interest, as it contains clues about the dominant dynamics of low-frequency flows (e.g., LeTraon et al., 2008; Xu and Fu, 2011). Because of the 10-day repeat track time, current state-of-the-art TOPEX/JASON satellite altimeters alias tides into longer periods (Parke et al., 1987). In contrast, because our model output is written at hourly intervals, we can easily separate low- from high-frequency signals in HYCOM. In regions such as the Kuroshio, where low-frequency motions are more energetic than tidal motions, the wavenumber spectrum of SSH is dominated by low-frequency motions. However, in regions where internal tides are energetic, for instance near Hawai'i, high-frequency motions dominate the high-wavenumber end of the wavenumber spectrum. The results of Richman et al. (2012) imply that internal tides will have to be accurately removed from data taken by the planned high-resolution wide-swath satellite altimeter (Fu et al., 2012) before low-frequency oceanic motions can be investigated. The accuracy with which internal tides can be removed from altimeter signals depends on the degree of non-stationarity of the internal tides. Internal tide non-stationarity in HYCOM is investigated in Shriver et al. (2014).




Figure 1. The internal tide amplitude of the principal lunar semidiurnal constituent M2 from the (a) altimetric-based and (b) HYCOM tidal analyses. The five subregions denoted by black boxes in (b) are used to compute area-averaged amplitudes in Shriver et al. (2012). From Shriver et al. (2012).

3. Insertion of topographic wave drag into eddying simulations

In Trossman et al. (2013) we again use the topographic wave drag parameterisation of Garner (2005), this time applied to low-frequency motions. The Trossman et al. (2013) simulations did not include tides. For simplicity, we refer to the Garner (2005) scheme here as a “wave drag” scheme even though it includes effects of topographic blocking, which yields low-level turbulence, as well as lee wave production and breaking. Maps of energy dissipation by quadratic bottom drag and by the parameterised topographic lee wave drag, both averaged over one year, are shown in Figure 2. As in the “offline” estimates of Nikurashin and Ferrari (2011) and Scott et al. (2011), as well as the offline estimates done in Trossman et al. (2013), the inline estimates shown in the bottom panel of Figure 2 indicate substantial lee wave drag dissipation in the Southern Ocean. The globally integrated internal lee wave drag dissipation in the inline Trossman et al. (2013) estimates is 0.4 TW, comparable to that seen in the offline estimates of Nikurashin and Ferrari (2011), Scott et al. (2011), and Trossman et al. (2013). However, as anticipated, there is indeed a strong feedback between the internal lee wave drag and the model flows. Adding wave drag to HYCOM substantially changes the modelled near-bottom eddy kinetic energy and stratification fields (Trossman et al., 2013).




Figure 2. Log10 of the bottom and wave drag terms in the energy budget of 1/12 eddying HYCOM simulation with , where each drag term has units [W m^{-2}]: (a) quadratic bottom boundary layer drag and (c) parameterised internal lee wave drag. Adapted from Trossman et al. (2013).

4. Ongoing and planned work

In ongoing and planned work with the HYCOM tide simulations, we are comparing the SSH frequency spectra in HYCOM versus tide gauges, as a measure of the accuracy of the partition of modelled low- versus high-frequency energy (Savage et al., paper in revision). We are estimating tidal aliasing errors in altimetrically-derived low-frequency wavenumber spectra and spectral fluxes (Savage et al., paper in preparation). We are also comparing the kinetic energy frequency spectra in HYCOM versus moored current meter records (Müller et al., paper in preparation). We are comparing both low- and high-frequency temperature variances (which are related to available potential energies) to temperature variances in moored historical records (Luecke et al. and Bassette et al., respective papers in preparation). To further elucidate the energetics of internal tides, we are preparing global maps of the baroclinic tidal energy fluxes and barotropic-to-baroclinic tidal energy conversions (Ansong et al. and Buijsman et al., papers in preparation). The latter work is part of the National Science Foundation-funded Climate Process Team project “Collaborative Research: Representing internal-wave driven mixing in global ocean models”, which focuses on improving estimates of internal-wave mediated mixing in the ocean, and is led by Professor Jennifer MacKinnon of the Scripps Institution of Oceanography. Finally, our long-term goals for the HYCOM tides work include improvement of tidal accuracy through the inclusion of data assimilation and better estimates of the self-attraction and loading term, and usage of the global HYCOM tidal solution to force regional and coastal models at their open-water boundaries.

In ongoing and planned work on parameterised topographic lee wave drag, we will investigate whether the inclusion of wave drag into eddying models improves the comparison of the models with observations. We will also compare the dissipation predictions of the Garner (2005) and Bell (1975) schemes with dissipation inferred from microstructure observations. Finally, we will investigate the impact of a more complete parameterisation of the vertical deposition of internal lee wave drag on the energetics and abyssal circulation in eddying models. These projects are all led by co-author Trossman (papers in preparation).

Acknowledgements

We thank Richard Ray for providing results from a global harmonic analysis of along-track satellite altimetry data, used in Figure 1. BKA, PGT and DST gratefully acknowledge support from National Science Foundation (NSF) grants OCE-0924481 and OCE-0960820, Naval Research Laboratory contract N000173-06-2-C003, and Office of Naval Research grants N00014-07-1-0392, N00014-09-1-1003 and N00014-11-1-0487. SRJ acknowledges NSF grant OCE-0960756. EPC acknowledges support from the Office of Naval Research grant N00014-09-1-0587. MCB, EJM, JGR, JFS, and AJW were supported by the project “Eddy resolving global ocean prediction including tides” sponsored by the Office of Naval Research under program element number 0602435N. Grants of computer time were provided by the Department of Defense (DoD) High Performance Computing Modernization Program. The simulations were performed on the SGI Altix Ice at the US Army Engineer Research and Development Center DoD Supercomputing Resource Center in Vicksburg, Mississippi. This is NRL contribution NRL/JA/7320-14-2170 and has been approved for public release.

References

- Arbic, B.K., S.T. Garner, R.W. Hallberg, and H.L. Simmons. 2004: The accuracy of surface elevations in forward global barotropic and baroclinic tide models. *Deep-Sea Research II*, 51, 3069-3101.
- Arbic, B.K., J.G. Richman, J.F. Shriver, P.G. Timko, E.J. Metzger, and A.J. Wallcraft, 2012: Global modeling of internal tides within an eddying ocean general circulation model. *Oceanography*, 25, 20-29, doi:10.5670/oceanog.2012.38.
- Arbic, B.K., J.F. Shriver, P.J. Hogan, H.E. Hurlburt, J.L. McClean, E.J. Metzger, R.B. Scott, A. Sen, O.M. Smedstad, and A.J. Wallcraft, 2009: Estimates of bottom flows and bottom boundary layer dissipation of the oceanic general circulation from global high-resolution models. *Journal of Geophysical Research*, 114, C02024.
- Arbic, B.K., A.J. Wallcraft and E.J. Metzger, 2010: Concurrent simulation of the eddying general circulation and tides in a global ocean model. *Ocean Modelling*, 32, 175-187, doi:10.1016/j.ocemod.2010.01.007.
- Bell, T.H., 1975: Lee waves in stratified flows with simple harmonic time dependence. *Journal of Fluid Dynamics*, 67, 705-722.
- Capet, X., J. C. McWilliams, M. J. Molemaker, and A. F. Shchepetkin, 2008: Mesoscale to submesoscale transition in the California current system. Part III: Energy balance and flux. *Journal of Physical Oceanography*, 38, 2256–2269.
- Chassignet, E.P., H.E. Hurlburt, E.J. Metzger, O.M. Smedstad, J. Cummings, G.R. Halliwell, R. Bleck, R. Baraille, A.J. Wallcraft, C. Lozano, H.L. Tolman, A. Srinivasan, S. Hankin, P. Cornillon, R. Weisberg, A. Barth, R. He, F. Werner, and J. Wilkin, 2009: U.S.

- GODAE: Global Ocean Prediction with the HYbrid Coordinate Ocean Model (HYCOM). *Oceanography*, 22(2), 64-75.
- Chassignet, E. P., J. G. Richman, E. J. Metzger, X. Xu, P. G. Hogan, B. K. Arbic, and A. J. Wallcraft, 2014: HYCOM high-resolution eddy simulations. *CLIVAR Exchanges*, 65, 22-25
- Egbert, G.D., A.F. Bennett, and M.G.G. Foreman, 1994: TOPEX/POSEIDON tides estimated using a global inverse model. *Journal of Geophysical Research*, 99, 24821-24852.
- Fu, L.-L., D. Alsdorf, R. Morrow, and E. Rodriguez, 2012: SWOT: The Surface Water and Ocean Topography Mission, Jet Propulsion Laboratory JPL-Publication 12-05, 228 pp
- Garner, S.T., 2005: A topographic drag closure built on an analytical base flux. *Journal of the Atmospheric Sciences*, 62, 2302-2315.
- Hecht, M.W., and H. Hasumi (Editors), 2008: Ocean modeling in an eddy regime. *Geophysical Monograph 177*, American Geophysical Union, Washington, 409 pp.
- Hendershott, M.C., 1972: The effects of solid earth deformation on global ocean tides. *Geophysical Journal of the Royal Astronomical Society*, 29, 389-402.
- McClean, J. L., A. J. Semtner, and V. Zlotnicki, 1997: Comparisons of mesoscale variability in the Semtner-Chervin quarter-degree model, the Los Alamos sixth-degree model, and TOPEX/POSEIDON Data. *Journal of Geophysical Research*, 102, 25203-25226.
- Le Traon, P.-Y., P. Klein, and B.-L. Hua, 2008: Do altimeter wavenumber spectra agree with the interior or surface quasigeostrophic theory? *Journal of Physical Oceanography*, 38, 1137-1142.
- Munk, W., and C. Wunsch, 1998: Abyssal recipes II: energetics of tidal and wind mixing. *Deep-Sea Research I*, 45, 1977-2010.
- Naveira-Garabato, A.C., K.L. Polzin, B.A. King, K.J. Heywood, and M. Visbeck, 2004: Widespread intense turbulent mixing in the Southern Ocean. *Science*, 303, 210-213.
- Nikurashin, M., and R. Ferrari, 2011: Global energy conversion rate from geostrophic flows into internal lee waves in the deep ocean. *Geophysical Research Letters*, 38, L08610.
- Parke, M.E., R.H. Stewart, D.L. Farless, and D.E. Cartwright, 1987: On the choice of orbits for an altimetric satellite to study ocean circulation and tides. *Journal of Geophysical Research*, 92, 11693-11707.
- Ray, R.D., 1998: Ocean self-attraction and loading in numerical tidal models. *Marine Geodesy*, 21, 181-191.
- Ray, R. D., and D. A. Byrne, 2010: Bottom pressure tides along a line in the southeast Atlantic Ocean and comparisons with satellite altimetry. *Ocean Dynamics*, 60, 1167-1176, doi:10.1007/s10236-010-0316-0.
- Richman, J.G., B.K. Arbic, J.F. Shriver, E.J. Metzger, and A.J. Wallcraft, 2012: Inferring dynamics from the wavenumber spectra of an eddy global ocean model with embedded tides. *Journal of Geophysical Research*, 117, C12012, doi:10.1029/2012JC008364.
- Scott, R.B., J.A. Goff, A.C. Naveira-Garabato, and A.J.G. Nurser, 2011: Global rate and spectral characteristics of internal gravity wave generation by geostrophic flow over topography. *Journal of Geophysical Research*, 116, C09029.
- Sen, A., R.B. Scott, and B.K. Arbic, 2008: Global energy dissipation rate of deep-ocean low-frequency flows by quadratic bottom boundary layer drag: Computations from current-meter data. *Geophysical Research Letters*, 35, L09606.
- Shriver, J.F., B.K. Arbic, J.G. Richman, R.D. Ray, E.J. Metzger, A.J. Wallcraft, and P.G. Timko, 2012: An evaluation of the barotropic and internal tides in a high resolution global ocean circulation model. *Journal of Geophysical Research*, 117, C10024, doi:10.1029/2012JC008170.
- Shriver, J.F., J.G. Richman, and B.K. Arbic, 2014: How stationary are the internal tides in a high resolution global ocean circulation model?, in press for *Journal of Geophysical Research Oceans*.
- Shum, C.K., P.L. Woodworth, O.B. Andersen, G.D. Egbert, O. Francis, C. King, S.M. Klosko, C. Le Provost, X. Li, J.-M. Molines, M.E. Parke, R.D. Ray, M.G. Schlax, D. Stammer, C.C. Tierney, P. Vincent, and C.I. Wunsch, 1997: Accuracy assessment of recent ocean tide models. *Journal of Geophysical Research*, 102, 25173-25194.
- Simmons, H.L., and M.H. Alford. 2012: Simulating the long range swell of internal waves generated by ocean storms, *Oceanography*, 25, 30-41, doi:10.5670/oceanog.2012.39.
- Simmons, H.L., R.W. Hallberg, and B.K. Arbic. 2004: Internal wave generation in a global baroclinic tide model. *Deep-Sea Research II*, 51, 3043-3068.
- Thoppil, P.G., J.G. Richman and P.J. Hogan, 2011: Energetics of a global ocean circulation model compared to observations. *Geophysical Research Letters*, 38, L15607, doi:10.1029/2011GL048347.
- Timko, P.G., B.K. Arbic, J.G. Richman, R.B. Scott, E.J. Metzger, and A.J. Wallcraft, 2012: Skill tests of three-dimensional tidal currents in a global ocean model: A look at the North Atlantic. *Journal of Geophysical Research*, 117, C08014, doi:10.1029/2011JC007617.
- Timko, P.G., B.K. Arbic, J.G. Richman, R.B. Scott, E.J. Metzger, and A.J. Wallcraft, 2013: Skill testing a three-dimensional global tide model to historical current meter records. *Journal of Geophysical Research Oceans*, 118, 6914-6933, doi:10.1002/2013JC009071.
- Trossman, D.S., B.K. Arbic, S.T. Garner, J.A. Goff, S.R. Jayne, E.J. Metzger, and A.J. Wallcraft, 2013: Impact of parameterized lee wave drag on the energy budget of an eddy global ocean model. *Ocean Modelling*, 72, doi:10.1016/j.ocemod.2013.08.006, 119-142.
- Wunsch, C., 1998: The work done by the wind on the oceanic general circulation. *Journal of Physical Oceanography*, 28, 2332-2340.
- Xu, Y., and L.-L. Fu., 2011: Global variability of the wavenumber spectrum of oceanic mesoscale turbulence. *Journal of Physical Oceanography*, 41, 802-809, doi:10.1175/2010JPO4558.1.